

HOCHSCHILD AND CYCLIC HOMOLOGY OF A FAMILY OF AUSLANDER ALGEBRAS

Rachel Taillefer*

Abstract

In this paper, we compute the Hochschild and cyclic homologies of the Auslander algebras of the Taft algebras. We also describe the first Chern character for the Taft algebras and for their Auslander algebras.

2000 Mathematics Subject Classification: 16E20, 16E40, 19D55.

Keywords: Hochschild homology, Cyclic homology, Auslander algebras, Chern characters.

1 Introduction

The object of this paper is to compute the Hochschild homology, the cyclic homology and the Chern characters of the Auslander algebras Γ_{Λ_n} of the Taft algebras Λ_n and of their Auslander algebras Γ_{Λ_n} , in order to study a possible influence of the Hopf algebra structure of Λ_n on them.

Note that Auslander algebras are useful when considering Artin algebras of finite representation type, since there is a bijection between the Morita equivalence classes of such algebras and the Morita equivalence classes of Auslander algebras (*cf.* [ARS]).

The Hopf algebra structure on an algebra Λ conveys an additional structure on the Grothendieck groups $K_0(\Lambda)$ and $\overline{K}_0(\Lambda)$ of isomorphism classes of projective (*resp.* all) indecomposable modules, since the tensor product over the base ring k of two Λ -modules is again a Λ -module, *via* the comultiplication of Λ . Furthermore, there is a one-to-one correspondence between the indecomposable modules over any algebra and the indecomposable projective modules over its Auslander algebra; in the case of a Hopf algebra, therefore, the Grothendieck group of projective modules of Γ_{Λ} is endowed with a multiplicative structure. However, this correspondence does not preserve the underlying vector spaces, and this multiplicative structure does not appear to be natural.

In this paper, we study the example of the Taft algebras; they are Hopf algebras which are neither commutative, nor cocommutative. They are interesting for various reasons; for instance, Λ_p is an example of a non-semisimple Hopf algebra whose dimension is the square of a prime (*cf.* [M]). They are of finite representation type; furthermore, when n is odd, Λ_n is isomorphic to the half-quantum group $u_q^+(\mathfrak{sl}_2)$ (q primitive n^{th} -root of unity), and is the only half-quantum group $u_q^+(\mathfrak{g})$ at a root of unity which is not of wild representation type (*cf.* [C1]). Then, for each n , Λ_n is not braided, but its Grothendieck group is a commutative ring nonetheless (*cf.* [C2, G]).

These examples show that the non-commutative, non-cocommutative Hopf algebra structure of Λ_n does not yield a natural multiplicative structure on its cyclic homology. There is a product, however, obtained by transporting that of $K_0(\Lambda)$ via the Chern characters, which are onto.

The paper is organised as follows: first, we recall the quiver of the Auslander algebras Γ_{Λ_n} and give a minimal projective resolution of Γ_{Λ_n} as a Γ_{Λ_n} -bimodule. Then, we compute the Hochschild and cyclic homologies of these algebras, and finally we compute the Chern characters of the algebras Λ_n and Γ_{Λ_n} .

Throughout this text, k is an algebraically closed field.


2 The quivers of the Auslander algebras Γ_{Λ_n}

In this paragraph, we shall describe the objects of our study: the Auslander algebras of the Taft algebras.

The Taft algebra Λ_n is described by quiver and relations as follows: the quiver is an oriented cycle with n vertices and n arrows, and the relations are all the paths of length greater than or equal to n .

*Supported by a Lavoisier grant from the French Ministry of Foreign Affairs.

Its Auslander algebra has been described in [GR] (see [ARS] p232 for a general definition). Its quiver is



where both vertical outer edges are identified (the quiver is on a cylinder). Let Q_n denote this quiver, let $\{e_{i,u}/(i,u) \in \mathbb{Z}/n\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}\}$ be the set of vertices of Q_n , and let $\{a_{i,u}; b_{i,u}/(i,u) \in \mathbb{Z}/n\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}\}$ be the set of edges of Q_n , as in the figure above.

The mesh relations on this quiver are: $a_{i,i-2}b_{i,i-1} = 0$ for all $i \in \mathbb{Z}/n\mathbb{Z}$ (the composition of two edges of any ‘triangle’ under the top diagonal is zero), and $a_{i,u-1}b_{i,u} + b_{i-1,u}a_{i,u} = 0$ for all i and u in $\mathbb{Z}/n\mathbb{Z}$ (the squares are anticommutative).

The algebra Γ_{Λ_n} is the quotient of the path algebra kQ_n by the ideal generated by these relations. We shall assume $n \geq 2$.

3 Hochschild and cyclic homologies of Γ_{Λ_n}

We are going to use the following theorem due to Happel to compute a minimal projective resolution of Γ_{Λ_n} as a Γ_{Λ_n} -bimodule (the situation in [H] is more general):

Theorem 3.1 ([H] 1.5) If

$$\dots \rightarrow R_p \rightarrow R_{p-1} \rightarrow \dots \rightarrow R_1 \rightarrow R_0 \rightarrow \Gamma_{\Lambda_n} \rightarrow 0$$

is a minimal projective resolution of Γ_{Λ_n} as a Γ_{Λ_n} -bimodule, then

$$R_p = \bigoplus_{\substack{(i,u) \\ (j,v)}} (\Gamma_{\Lambda_n} e_{j,v} \otimes e_{i,u} \Gamma_{\Lambda_n})^{\dim_k \text{Ext}_{\Gamma_{\Lambda_n}}^p(S_{i,u}; S_{j,v})}.$$

In our case, we have:

Corollary 3.2 The complex

$$\begin{aligned} \dots 0 \rightarrow 0 \rightarrow \bigoplus_{\substack{(i,u) \\ i \neq u}} \Gamma_{\Lambda_n} e_{i-1,u-1} \otimes e_{i,u} \Gamma_{\Lambda_n} &\rightarrow \bigoplus_{(i,u)} [(\Gamma_{\Lambda_n} e_{i-1,u} \otimes e_{i,u} \Gamma_{\Lambda_n}) \oplus (\Gamma_{\Lambda_n} e_{i,u-1} \otimes e_{i,u} \Gamma_{\Lambda_n})] \\ &\rightarrow \bigoplus_{(i,u)} \Gamma_{\Lambda_n} e_{i,u} \otimes e_{i,u} \Gamma_{\Lambda_n} \rightarrow \Gamma_{\Lambda_n} \rightarrow 0 \end{aligned}$$

is a minimal projective resolution of Γ_{Λ_n} as a Γ_{Λ_n} -bimodule.

Proof: We need to compute the Ext groups between the simple Γ_{Λ_n} -modules. First, let us compute the projective resolutions of the simple modules:

Lemma 3.3 Let $P_{i,u}$ denote the indecomposable projective Γ_{Λ_n} -module at the vertex $e_{i,u}$, and let $S_{i,u} = \text{top}(P_{i,u})$ be the corresponding simple module. The minimal projective resolutions of the simple modules are:

$$\begin{array}{ccccccccccc} 0 & \longrightarrow & P_{i-1,i-2} & \longrightarrow & P_{i-1,i-1} & \longrightarrow & P_{i,i} & \longrightarrow & S_{i,i} & \longrightarrow & 0 \\ 0 & \longrightarrow & P_{i-1,i-2} & \longrightarrow & P_{i,i-2} & \longrightarrow & P_{i,i-1} & \longrightarrow & S_{i,i-1} & \longrightarrow & 0 \\ 0 & \longrightarrow & P_{i-1,i-j-1} & \longrightarrow & P_{i-1,i-j} \oplus P_{i,i-j-1} & \longrightarrow & P_{i,i-j} & \longrightarrow & S_{i,i-j} & \longrightarrow & 0 \end{array}$$

for $2 \leq j \leq n-1$.

Proof of Lemma: We consider only the $S_{n-1,u}$, because the other cases may be obtained by translating the quiver along the cylinder on which it lies. It is then straightforward to compute their minimal projective resolutions. $\square$

We can now compute the Ext groups:

Lemma 3.4 Let S be a simple Γ_{Λ_n} -module. Then:

$$\begin{aligned} \text{Ext}_{\Gamma_{\Lambda_n}}^0(S_{i,u}; S) &= \begin{cases} k & \text{if } S = S_{i,u}, \\ 0 & \text{if } S \neq S_{i,u}, \end{cases} \\ \text{Ext}_{\Gamma_{\Lambda_n}}^1(S_{i,i}; S) &= \begin{cases} k & \text{if } S = S_{i-1,i}, \\ 0 & \text{if } S \neq S_{i-1,i}, \end{cases} \\ \text{Ext}_{\Gamma_{\Lambda_n}}^1(S_{i,i-1}; S) &= \begin{cases} k & \text{if } S = S_{i,i-2}, \\ 0 & \text{if } S \neq S_{i,i-2}, \end{cases} \\ \text{Ext}_{\Gamma_{\Lambda_n}}^1(S_{i,i-j}; S) &= \begin{cases} k & \text{if } S = S_{i,i-j-1} \text{ or } S = S_{i-1,i-j}, \\ 0 & \text{if } S \neq S_{i-1,i-j-1} \text{ and } S \neq S_{i-1,i-j}, \end{cases} \quad \text{if } 1 \leq j \leq n-1 \\ \text{Ext}_{\Gamma_{\Lambda_n}}^2(S_{i,i}; S) &= 0 \\ \text{Ext}_{\Gamma_{\Lambda_n}}^2(S_{i,i-j}; S) &= \begin{cases} k & \text{if } S = S_{i-1,i-j-1}, \\ 0 & \text{if } S \neq S_{i-1,i-j-1}, \end{cases} \quad \text{if } 1 \leq j \leq n-1 \\ \text{Ext}_{\Gamma_{\Lambda_n}}^p(S_{i,u}; S) &= 0 \text{ if } p \geq 3. \end{aligned}$$

Applying Happel's Theorem (3.1) we get the minimal projective resolution for Γ_{Λ_n} . $\blacksquare$

Now applying the functor $\Gamma_{\Lambda_n} \otimes_{\Gamma_{\Lambda_n} - \Gamma_{\Lambda_n}} ?$ to this resolution, we obtain a complex:

$$\dots 0 \longrightarrow \dots \longrightarrow 0 \longrightarrow kQ_0 \longrightarrow 0.$$

Therefore:

Proposition 3.5 The Hochschild homology of Γ_{Λ_n} is:

$$\begin{cases} \text{HH}_0(\Gamma_{\Lambda_n}) = kQ_0 \cong k^{n^2} \\ \text{HH}_p(\Gamma_{\Lambda_n}) = 0 \end{cases} \quad \forall p > 0,$$

and hence the cyclic homology of Γ_{Λ_n} is:

$$\begin{cases} \text{HC}_{2p}(\Gamma_{\Lambda_n}) = kQ_0 \cong k^{n^2} \\ \text{HC}_{2p+1}(\Gamma_{\Lambda_n}) = 0 \end{cases} \quad \forall p \geq 0.$$

Remark 3.6 There doesn't seem to be any connection between these results and those for Λ_n . Indeed, the Hochschild and cyclic homologies for the Taft algebras are given as follows (see [S] for the Hochschild homology and [T, T1] for the cyclic homology):

$$\begin{cases} \text{HH}_0(\Lambda_n) = k^n \\ \text{HH}_p(\Lambda_n) = k^{n-1} \end{cases} \quad \forall p > 0.$$

and

$$\begin{cases} \text{HC}_{2p}(\Lambda_n) = k^n, \\ \text{HC}_{2p+1}(\Lambda_n) = k^{n-1} \end{cases} \quad \forall p \geq 0.$$

4 Chern characters of Λ_n and Γ_{Λ_n}

Let $K_0(\Lambda_n)$ (*resp.* $K_0(\Gamma_{\Lambda_n})$) be the Grothendieck group of projective Λ_n -modules (*resp.* Γ_{Λ_n} -modules). We are interested in the Chern characters $ch_{0,p} : K_0(\Lambda_n) \rightarrow HC_{2p}(\Lambda_n)$ (*resp.* $K_0(\Gamma_{\Lambda_n}) \rightarrow HC_{2p}(\Gamma_{\Lambda_n})$). We shall write $[P_j]$ (*resp.* $[P_{i,u}]$) for the isomorphism class of the projective module at the vertex e_j (*resp.* $e_{i,u}$).

Set $\sigma^p = (y_p, z_p, \dots, y_1, z_1, y_0) \in \mathbb{N}^{2p+1}$ with $y_p = (-1)^p(2p)!/p!$ and $z_p = (-1)^{p-1}(2p)!/2(p!)$. There is a system of generators of $HC_{2p}(\Lambda_n)$ (*resp.* $HC_{2p}(\Gamma_{\Lambda_n})$) given by the following set:

$$\{\sigma_i^p := \sigma^p(e_i, \dots, e_i) \in (\text{Tot } CC(\Lambda_n))_{2p} / i = 0, \dots, n-1\}$$

(*resp.* by $\{\sigma_{i,u}^p := \sigma^p(e_{i,u}, \dots, e_{i,u}) \in (\text{Tot } CC(\Gamma_{\Lambda_n}))_{2p} / i, u \in \{0, 1, \dots, n-1\}\}$).

Consider the elements

$$\begin{array}{ccc} \epsilon_j : \Lambda_n & \longrightarrow & \Lambda_n \\ \lambda & \mapsto & \lambda e_j \end{array} \quad \text{and} \quad \begin{array}{ccc} \epsilon_{i,u} : \Gamma_{\Lambda_n} & \longrightarrow & \Gamma_{\Lambda_n} \\ \lambda & \mapsto & \lambda e_{i,u} \end{array}$$

in $\mathcal{M}_1(\Lambda_n)$ and $\mathcal{M}_1(\Gamma_{\Lambda_n})$; their ranges are the corresponding projective modules. Then by definition of the Chern characters (see [L] 8.3.4), we have:

$$\begin{aligned} ch_{0,p}([P_j]) &= ch_{0,p}([\epsilon_j]) := \text{tr}(c(\epsilon_j)) = \sigma_j^p \text{ in } HC_{2p}(\Lambda_n) \\ ch_{0,p}([P_{i,u}]) &= ch_{0,p}([\epsilon_{i,u}]) = \sigma_{i,u}^p \text{ in } HC_{2p}(\Gamma_{\Lambda_n}). \end{aligned}$$

using the isomorphisms $\mathcal{M}_m(\Lambda) \cong \mathcal{M}_m(k) \otimes \Lambda$. Here,

$$c(\epsilon_j) = (y_p \epsilon_j^{\otimes 2p+1}, z_p \epsilon_j^{\otimes 2p}, \dots, z_1 \epsilon_j^{\otimes 2}, y_0 \epsilon_j) \in \mathcal{M}(\Gamma_{\Lambda_n})^{\otimes 2p+1} \oplus \dots \oplus \mathcal{M}(\Gamma_{\Lambda_n}).$$

Remark 4.1 There is a decomposition formula for the tensor product of indecomposable modules on Λ_n (see [C2, G]). From this formula, we get inductively:

$$ch_{0,p}([L_1] \otimes \dots \otimes [L_r]) = \frac{1}{n^2} \prod_{i=1}^r (\dim L_i) (\sigma_0^p, \dots, \sigma_{n-1}^p), \text{ for } r \geq 2,$$

where the L_i are arbitrary projective Λ_n -modules. Unfortunately, this product in the cyclic homology doesn't seem natural.

Remark 4.2 Let $\overline{K}_0(\Lambda_n)$ be the Grothendieck group of all Λ_n -modules (not just the projective ones). Then $\overline{K}_0(\Lambda_n) \cong K_0(\Gamma_{\Lambda_n})$. Hence, if $N_{i,u}$ is the indecomposable Λ_n -module which starts at the vertex i and ends at the vertex u , it corresponds to the projective Γ_{Λ_n} -module $P_{i,u}$, and we get a map:

$$\begin{array}{ccc} \overline{K}_0(\Lambda_n) & \longrightarrow & HC_{2p}(\Gamma_{\Lambda_n}) \\ N_{i,u} & \mapsto & \sigma_{i,u}^p. \end{array}$$

Remark 4.3 Although Γ_{Λ_n} is not a Hopf algebra, its Grothendieck group $K_0(\Gamma_{\Lambda_n})$ does have a ring structure, which does not appear to be natural: for every $[P]$ in $K_0(\Gamma_{\Lambda_n})$, there exists a $[B]$ in $K_0(\Lambda_n)$ such that $[P] = [\text{Hom}_{\Lambda_n}(M, B)]$, where M is the sum of all isomorphism classes of indecomposable Λ_n -modules. If $[Q] = [\text{Hom}_{\Lambda_n}(M, C)]$ is another element in $K_0(\Gamma_{\Lambda_n})$, we can set

$$[P] \cdot [Q] = [\text{Hom}_{\Lambda_n}(M, B \otimes_k C)]$$

(the vector space $B \otimes_k C$ is a Λ_n -module since Λ_n is a Hopf algebra). In fact, using the decomposition in [C2, G], the product can be written:

$$[P_{i,u}][P_{j,v}] = \begin{cases} \sum_{l=0}^{v-j} [P_{i+j+l, u+v-l}] & \text{if } u+v-(i+j) \leq n-1 \\ \sum_{l=0}^e [P_{i+j+l, u+v+l-1}] + \sum_{m=e+1}^{v-j} [P_{i+j+m, u+v-m}] & \text{if } e := u+v-(i+j)-(n-1) \geq 0 \end{cases}$$

where u and v represent elements in $\mathbb{Z}/n\mathbb{Z}$ such that $u-i$ and $v-j$ are in $\{0, 1, \dots, n-1\}$.

References

- [ARS] M. AUSLANDER, I. REITEN AND S.O. SMALØ, Representation Theory of Artin Algebras, *Cambridge Studies in Advanced Mathematics*, 36. Cambridge University Press, Cambridge, (1997).
- [C1] C. CIBILS, Half-quantum Groups at Roots of Unity, Path Algebras, and Representation Type, *Internat. Math. Res. Notices* (1997), no. 12, 541–553.
- [C2] C. CIBILS, A Quiver Quantum Group, *Comm. Math. Phys.* **157** (1993), no. 3, pp 459-477.
- [GR] P. GABRIEL AND CH. RIEDTMANN, Group Representations Without Groups, *Comment. Math. Helv.* **54** (1979), no. 2, pp 240-287.
- [G] E. GUNNLAUGSDÓTTIR, Monoidal Structure of the Category of u_q^+ -modules, *to appear in Linear Algebra and its Applications*.
- [H] D. HAPPEL, Hochschild Cohomology of Finite-dimensional Algebras, *Séminaire d'Algèbre Paul Dubreil et Marie-Paul Malliavin, 39ème Année (Paris, 1987/1988)*, pp 108-126, *Lecture Notes in Math.* **1404**, Springer, Berlin-New York (1989).
- [L] J-L. LODAY, Cyclic Homology, Appendix E by María O. Ronco, *Springer-Verlag, Berlin*, (1992).
- [M] S. MONTGOMERY, Classifying Finite-dimensional Semisimple Hopf Algebras, *Contemp. Math.* **229** (1998), pp 265-279, *Amer. Math. Soc., Providence, RI*.
- [S] E. SKÖLDBERG, The Hochschild Homology of Truncated and Quadratic Monomial Algebras, *J. London Math. Soc. (2)* **59** (1999), no. 1, pp 76-86 (1999).
- [T] R. TAILLEFER, Théories Homologiques des Algèbres de Hopf, *Thèse de Doctorat de l'Université Montpellier II* [with an abridged version in English] (2001).
- [T1] R. TAILLEFER, Cyclic Homology of Hopf Algebras, *K-Theory* **24**, pp 69-85 (2001).

RACHEL TAILLEFER
 DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE,
 UNIVERSITY OF LEICESTER,
 LEICESTER LE1 7RH,
 UNITED KINGDOM.
 E-MAIL: R.Taillefer@mcs.le.ac.uk.